

Effectiveness of Nurse Led Intervention on Respiratory Parameters and Cholinesterase among Organophosphorus Compound Poisoning Patients Admitted in Dhanvantri Critical Care Centre at Erode – A Historical Comparison

Sampoornam. W, Aruna Devi. G, Karthika. R, Mariya pushpam. S, Pavithra. R, Surya. M, Saranya. R, Gnana Ashwini. A

Reader, Mental health nursing department, Dhanvantri College of Nursing Pallakkapalayam, Namakkal
IV Year Bsc (N) Students, Dhanvantri College of Nursing, Pallakkapalayam.

ARTICLE INFO

Article history:

Published online: 30th Sept, 2017

Key words:

Nurse Led Intervention

Respiratory Parameters

Cholinesterase

Organophosphorus Compound
Poisoning

ABSTRACT

Context: Nurses play a vital role in the management of poisoning, as it demands close observation, timely administration of antidotes in adequate doses and skilful nursing interventions. **Methods:** Quasi experimental design- A historical comparison was adopted. Total 30 organophosphorus compound poisoning patients, out of which 15 participants were in experimental group and 15 participants were in control group. Purposive sampling technique was used to select the participants admitted in Dhanvantri Critical Care Centre at Erode. The level of respiratory parameters were assessed by adopting in vitro ABG (Arterial Blood Gas) analysis which constitutes pH (acidity), pCO₂ (partial pressure of carbon dioxide), pO₂ (partial pressure of oxygen) and HCO₃ (bicarbonate). Serum cholinesterase estimation level was measured from the vein by adopting in vitro method. **Results:** The Paired t' test value of respiratory parameter (pH, pCO₂, pO₂, HCO₃) scores showed statistical significance in experimental group. Only pCO₂ paired t' test scores showed statistical significance in control group. Unpaired t' test value of posttest respiratory parameter (pH, pCO₂, pO₂, HCO₃) scores between experimental and control group was statistically significant. The Paired t' test value of cholinesterase scores showed statistical significance in experimental group (t=4.00). Whereas control group showed statistical non significance (t=0.71). Unpaired t' test value of posttest cholinesterase scores between experimental and control group was statistically significant (t=3.84). **Conclusion:** The study findings revealed that administration of Nurse Led Intervention were highly significant to improve respiratory parameters and levels of cholinesterase among organophosphorus compound poisoning patients in experimental group than control group.

Introduction

Globally, organophosphorus (OP) pesticide poisoning is a serious occupational hazard accounting for more than

80% of pesticide related hospitalisation. Pesticide self poisoning is responsible in killing approximately 300,000 people worldwide every year and mostly from rural background. In developing countries the mortality can be

as high as 70% (Eddleston M. 2000, Eddleston M, Phillips MR. 2004). High mortality could be probably due to lack of hospital services in the vicinity, inadequate transport facility, more patients with care givers ratio and finally non availability of definite antidote (Buckley NA, 2004).

Recent data from National crime bureau of India shows suicide by consumption of pesticides account for 19.4% and 19.7% of all cases of suicidal poisoning in the year 2006 and 2007 respectively. WHO estimates that approximately 3 million pesticide poisoning occur worldwide and cause more than 2,20,000 deaths. Developing countries like India and Srilanka have reported increased rates of toxicity and death (Darren M 2007).

Respiratory complications of OP poisoning occur during, and as a consequence of, the acute cholinergic crisis, delayed neuromuscular dysfunction, and recurrent cholinergic toxicity. Patients who were intubated after 24 hours required a significantly longer period of ventilation. Bronchorrhea results from neuronal and nonneuronal cholinergic stimulation of the mucus glands, cilia, and cells producing periciliary fluid (Kummer W, Lips KS, Pfeil U 2008, Takemura Y, Helms MN et al, 2013). Acute respiratory distress syndrome (ARDS) has been reported in several cases of OP insecticide poisoning (Akur S, Veral A, Ege B, 2008). It may result directly from pulmonary complications of poisoning, such as aspiration or inhalation, or indirectly via hematogenous exposure to OP compounds.

It was observed that majority of the cases were showing respiratory alkalosis (65%) on ABG, followed by respiratory acidosis (15%) and metabolic alkalosis (5%). No cases showed metabolic acidosis in the study. 15% cases showed normal acid base status in the study. The cases of respiratory acidosis and apodosis were further classified in to acute, partly compensated and fully compensated. Out of the total 20 cases 2 cases (10%) showed acute respiratory acidosis whereas 1 case (5%) showed partly compensated respiratory acidosis. 4 cases (20%) showed acute respiratory alkalosis. Whereas, partly and fully compensated respiratory alkalosis was observed in 6 (30%) and 3 (15%) cases respectively (Dudekula Moulali, D.V.H.S Sarma, et al, 2014).

Clinical manifestations following OP poisoning can be associated with the extent of decrease of PChE. OP pesticides inhibit carboxylic esterase enzymes including acetylcholinesterase (AChE) and plasma cholinesterase (PChE). AChE can be found in erythrocytes, nervous tissue and skeletal muscles, while PChE can be found in plasma, liver, heart, pancreas and brain. Most of clinical manifestations associated with exposure to OP compounds have been attributed to inhibition of these enzymes (Karalliedde L, 2002).

Organophosphate poisoning is a common cause of acute poisoning in India with high mortality. Prompt recognition and aggressive treatment of acute intoxication is essential to minimize the mortality and morbidity.

Nurses play a vital role in the management of poisoning, as it demands close observation, timely administration of antidotes in adequate doses and skilful nursing interventions. Counselling to the poisoned patients will reduce the chances of a repeat attempt at poisoning. It also enables the health care personnel to improve the quality of treatment, minimize the cost of therapy and the period of hospitalization. Family counseling is mandated; this helps the family members to cope with the situation and accept the patient as he is.

Poisoning is a common cause of hospital admissions. Caring of patients with OP poisoning is a challenge for nurses. Assessment and prevention of complications is one of the vital roles of the nurse. Appropriate, evidenced based practice will enhance quick recovery, reduce morbidity and mortality. Nurses must ensure that both patients and family members receive counselling, to cope and live in the community (Amala Rajan, Ilavarasi Jesudoss, 2016). Many literatures shows paucity of research work in nurse led intervention on respiratory parameters and cholinesterase among organophosphorus compound poisoning patients. Still research gap exists with nurse led intervention for respiratory parameters and cholinesterase in organophosphorus compound poisoning patients. The novelty of this research work is based on the respiratory parameters and cholinesterase level and need for nurse led intervention in which control group participants were selected from the previous case reported within 6 months.

1 Materials and Methods

Prior to the collection of data, permission was obtained from the Chairman, Dhanvantri Critical Care Centre at Erode. This study was approved by Institutional Human Ethical Committee. After obtaining written informed consent, organophosphorus compound poisoning patients who fulfilled the inclusion criteria were recruited and enrolled in the study on June 2016. Quasi experimental design- A historical comparison was adopted. Total 30 organophosphorus compound poisoning patients, out of which 15 participants were in experimental group and 15 participants were in control group. Purposive sampling technique was used to select the participants admitted in Dhanvantri Critical Care Centre at Erode.

All participants were interviewed before intervention in experimental and control group by assessing demographic variables like age, gender, occupation, family monthly income, residential area, intention of poison and type of poison. The level of respiratory parameters were assessed by adopting in vitro ABG (Arterial Blood Gas) analysis which constitutes pH (acidity), pCO₂ (partial pressure of carbon dioxide), pO₂ (partial pressure of oxygen) and HCO₃ (bicarbonate). Serum cholinesterase estimation level was measured from the vein by adopting in vitro method. For both the test, blood sample was collected during morning around 4-5 a.m. ABG (Arterial Blood Gas) analysis was done daily, whereas serum cholinesterase estimation was done once in 5 days. 2 mL

of arterial blood was collected from femoral artery and 2 mL of venous blood was collected from antecubital fossa and transferred to the EDTA vacutainer. It was labeled with patient's name, age and serial number. Throughout the procedure aseptic measures were taken. The collected blood was sent to Dhanvantri Critical Care laboratory by following standard operating procedure.

Immediately after pretest the participants received Nurse Led Intervention protocol by administering high flow of oxygen (5-7 litres) based on oxygen saturation with subsequent frequent suctioning once in 2 hours and brief psychotherapy focusing to raise awareness of problem and explores problem solving techniques on primarily present concern and stressors once a day. Intervention was given on 1st day after weaning from the ventilator for 3 consecutive days. Control group participants were selected from the previous case reported within 6 months and compared with experimental subjects.

Subject's right to withdraw/withhold the information was

ensured before data collection. Informed consent form was translated into Tamil language. After clarifying the doubts, signature was obtained from organophosphorus compound poisoning patients. Written Informed consent was obtained from each participants related to the study purpose, type of data, nature of commitments, participations and procedure. Confidentiality of the data was ensured throughout the study. Risk benefit ratio was calculated.

2 Results

Descriptive, inferential and nonparametric statistical analytic method was used to analyze and interpret the data. A probability of 0.05 or less was taken as statistically significant. Statistical package for social science, PCT version 17 (SPSS Inc, Chicago) was used for analysing the data.

Table 1 Frequency and Percentage distribution of pre and posttest scores on respiratory parameters in experimental group

| Respiratory Parameters | Levels of Respiratory Acidosis | Experimental group | | | | Levels of Respiratory Alkalosis | Experimental group | | | |
|---|--------------------------------|--------------------|-----|----------|-----|---------------------------------|--------------------|-----|----------|-----|
| | | Pretest | | Posttest | | | Pretest | | Posttest | |
| | | F | % | F | % | | F | % | F | % |
| pH pCO ₂ pO ₂ HCO ₃ | Mild | 4 | 27% | 6 | 40% | Mild | 6 | 40% | 3 | 21% |
| | Moderate | 2 | 13% | 2 | 13% | Moderate | 1 | 7% | 2 | 13% |
| | Severe | 1 | 7% | 2 | 13% | Severe | 2 | 13% | - | - |
| n = 15 | | | | | | | | | | |

In pretest, experimental group showed 27% with mild respiratory acidosis, 13% with moderate respiratory acidosis and 7% with severe respiratory acidosis. In posttest, 40% had mild respiratory acidosis, 13% of participants showed moderate respiratory acidosis and 13% showed severe respiratory acidosis. Likewise in

pretest, experimental group showed 40% with mild respiratory alkalosis, 7% with moderate respiratory alkalosis and 13% with severe respiratory alkalosis. In posttest, 21% had mild respiratory alkalosis, 13% of participants showed moderate respiratory alkalosis and none showed severe respiratory alkalosis (Table 1).

Table 2 Frequency and Percentage distribution of pre and posttest scores on respiratory parameters in control group

| Respiratory Parameters | Levels of Respiratory Acidosis | Control group | | | | Levels of Respiratory Alkalosis | Control group | | | |
|---|--------------------------------|---------------|-----|----------|-----|---------------------------------|---------------|-----|----------|-----|
| | | Pretest | | Posttest | | | Pretest | | Posttest | |
| | | F | % | F | % | | F | % | F | % |
| pH pCO ₂ pO ₂ HCO ₃ | Mild | 5 | 33% | 4 | 27% | Mild | 2 | 13% | 1 | 7% |
| | Moderate | 4 | 27% | 2 | 13% | Moderate | 1 | 7% | 6 | 40% |
| | Severe | 2 | 13% | - | - | Severe | 1 | 7% | 2 | 13% |
| n = 15 | | | | | | | | | | |

In pretest, control group showed 33% with mild respiratory acidosis, 27% with moderate respiratory acidosis and 13% with severe respiratory acidosis. In posttest, 27% had mild respiratory acidosis, 13% of participants showed moderate respiratory acidosis and none showed severe level of respiratory acidosis. Likewise in pretest, control group showed 13% with mild

respiratory alkalosis, 7% with moderate respiratory alkalosis and 7% with severe respiratory alkalosis. In posttest, 7% had mild respiratory alkalosis, 40% of participants showed moderate respiratory alkalosis and 13% showed severe level of respiratory alkalosis (Table 2).

In experimental group, the measured cholinesterase level

in pretest showed low elevation (53%), 27% participants reported with average elevation and 20% revealed high elevation of cholinesterase. In posttest 20% had low elevation of cholinesterase, 20% showed average elevation and 60% showed high elevation of cholinesterase. Control group in pretest showed 60%

low elevation, 20% average elevation and 20% high elevation of cholinesterase. In posttest 40% had low elevation of cholinesterase, 13% showed average elevation and 47% showed high elevation of cholinesterase (Table 3).

Table 3 Frequency and percentage distribution of pre and posttest scores on cholinesterase level among organophosphorus compound poisoning patients in experimental group and control group

| Level of Cholinesterase | Experimental group | | | | Control group | | | |
|-------------------------|--------------------|-----|----------|-----|---------------|-----|----------|-----|
| | Pretest | | Posttest | | Pretest | | Posttest | |
| | F | % | F | % | F | % | F | % |
| Low | 8 | 53% | 3 | 20% | 9 | 60% | 6 | 40% |
| Average | 4 | 27% | 3 | 20% | 3 | 20% | 2 | 13% |
| High | 3 | 20% | 9 | 60% | 3 | 20% | 7 | 47% |
| N =30 (n1=15, n2=15) | | | | | | | | |

Table 4 Comparison of mean, standard error, Paired t' test and Unpaired t' test value with pre and posttest scores of respiratory parameters between experimental and control group

| Variable | Group | Mean ± SE | Group | Mean ± SE | Significance Paired 't' test | | Significance Unpaired 't' test | | | |
|------------------------|----------------------|-----------|--------------------|-------------|------------------------------|----------------------|--------------------------------|---------------|------|--|
| | | | | | Experimental group | Control group | Experimental group | Control group | | |
| Respiratory parameters | pH | Pretest | Experimental group | 2.87 ± 0.09 | Control group | t = 3.50 P<0.003 | t = 1.87 P >0.08 | Mean | Mean | |
| | | Posttest | | 2.40 ± 0.16 | | | | 2.60 | 1.93 | |
| | pCO2 | Pretest | Experimental group | 1.80 ± 0.20 | Control group | t = 6.50 P <0.001 | t = 2.25 P <0.04 | Mean | Mean | |
| | | Posttest | | 2.07 ± 0.25 | | | | 1.73 | 1.20 | |
| | pO2 | Pretest | Experimental group | 2.47 ± 0.19 | Control group | t = 2.64 P <0.01 | t = 1.46 P >0.16 | Mean | Mean | |
| | | Posttest | | 2.13 ± 0.26 | | | | 2.13 | 1.33 | |
| | HCO3 | Pretest | Experimental group | 2.00 ± 0.22 | Control group | t = 2.64 P <0.01 | t = 1.46 P >0.16 | Mean | Mean | |
| | | Posttest | | 2.33 ± 0.21 | | | | 2.33 | 1.80 | |
| | N =30 (n1=15, n2=15) | | | | | | | | | |

Comparison of mean, standard error of pretest pH scores in experimental group showed 2.87±0.09 and posttest

showed 2.40 ± 0.16 . Pretest pCO₂ mean scores in experimental group showed 1.80 ± 0.20 and posttest showed 2.07 ± 0.25 . Pretest pO₂ mean scores in experimental group showed 2.47 ± 0.19 and posttest showed 2.13 ± 0.26 . Pretest HCO₃ mean scores in experimental group showed 2.00 ± 0.22 and posttest showed 2.33 ± 0.21 (Table 4)

Likewise, comparison of mean, standard error of pretest pH scores in control group showed 2.27 ± 0.21 and posttest showed 2.07 ± 0.23 . Pretest pCO₂ mean scores in control group showed 1.80 ± 0.14 and posttest showed 2.67 ± 0.13 . Pretest pO₂ mean scores in control group showed 2.13 ± 0.19 and posttest showed 2.00 ± 0.17 . Pretest HCO₃ mean scores in control group showed 1.93 ± 0.21 and posttest showed 2.07 ± 0.21 (Table 4)

The Paired t' test value of respiratory parameter (pH, pCO₂, pO₂, HCO₃) scores presented in Table-4 showed statistical significance in experimental group. Whereas, Paired t' test value of respiratory parameter (pH, pO₂, HCO₃) scores showed statistical non significance in control group. Only pCO₂ paired t' test scores showed statistical significance in control group (Table 4)

Unpaired t' test value of posttest respiratory parameter (pH, pCO₂, pO₂, HCO₃) scores between experimental and control group was statistically significant. It indicates that there was significant difference between the groups. Based on posttest pH, pCO₂, pO₂, HCO₃ mean scores, experimental group showed improvement in all respiratory parameters compared with control group (Table 4)

Table 5 Comparison of mean, standard error, Paired t' test and Unpaired t' test value with pre and posttest scores of cholinesterase level between experimental and control group

| Variable | Group | Mean ± SE | Significance Paired 't' test | | Significance Unpaired 't' test | |
|-----------------------|----------------------------|-----------------|---|---|-----------------------------------|------------------------------|
| | | | Experimental group Pre – Posttest | Control Group Pre – Posttest t | Experimental group Posttest | Control group Posttest |
| Cholinesterase | Experimental – Pretest | 1.80 ± 0.22 | t = 4.00 P < 0.001 | t = 0.71 P > 0.48 | Mean= 2.13 | Mean= 1.27 |
| | Experimental – Posttest | 1.27 ± 0.12 | | | t = 3.84 P < 0.001 | |
| | Control- Pretest | 1.80 ± 0.20 | | | | |
| | Control - Posttest | 1.53 ± 0.19 | | | | |
| N = 30 (n1=15, n2=15) | | | | | | |

Comparison of mean, standard error of pretest cholinesterase scores in experimental group showed 1.80 ± 0.22 and posttest showed 1.27 ± 0.12 . Likewise, comparison of mean, standard error of pretest cholinesterase scores in control group showed 1.80 ± 0.20 and posttest showed 1.53 ± 0.19 (Table 5).

The Paired t' test value of cholinesterase scores presented in (Table-5) showed statistical significance in experimental group (t=4.00). Whereas control group showed statistical non significance (t=0.71).

Unpaired t' test value of posttest cholinesterase scores between experimental and control group was statistically significant (t=3.84). It indicates that there was significant difference between the groups. Based on posttest cholinesterase mean scores, experimental group (Mean=2.13) showed increased level of cholinesterase compared with control group (Mean=1.27).

3 Discussion

Distribution of demographic variable in experimental group according to their age group depicts that highest percentage (67%) were in the age group of 31-40 years. In control group according to their age group highest percentage (40%) were in the age group of 31-40 years. Gender depicts that highest percentage (53%) were female gender in experimental group. In control group highest percentages (73%) were male gender. This result is supported with male to female ratio of 1:1.5. The mean age of the patients was 25.5 with a standard deviation of 9.45 (Getnet Mequanint Adinew & Assefa Belay Asrie, 2017). Participants according to their residential area showed majority 60% were from rural area in experimental group and in control group majority 67% were from rural area. A Retrospective study showed that 56.7% of the cases lived in an urban environment compared to 43.3% who lived rurally (Getnet Mequanint Adinew & Assefa Belay Asrie, 2017). Type of poisoning depicts that highest 53% of participants had taken pesticide poisoning in experimental

group. Likewise highest 67% of participants had taken insecticide poisoning in control group. Significant result reported by Subhash Chandra Joshi & Chandra Prakash, (2013) revealed that the most common motive of poisoning was with a suicidal intent, both in males (50.80%) and females (43.01%) and highest number of patients consumed Dichlorvos (40.86%).

Nurses play a vital role in the management of poisoning, as it demands close observation, timely administration of antidotes in adequate doses and skilful nursing interventions. Poisoning is a common cause of hospital admissions. Caring of patients with OP poisoning is a challenge for nurses. Assessment and prevention of complications is one of the vital roles of the nurse. Appropriate, evidenced based practice will enhance quick recovery, reduce morbidity and mortality. Nurses must ensure that both patients and family members receive counselling, to cope and live in the community (Amala Rajan, Ilavarasi Jesudoss, 2016). The Paired t' test value of respiratory parameter (pH, pCO₂, pO₂, HCO₃) scores showed statistical significance in experimental group. Whereas, Paired t' test value of respiratory parameter (pH, pO₂, HCO₃) scores showed statistical non significance in control group. Only pCO₂ paired t' test scores showed statistical significance in control group. Unpaired t' test value of posttest respiratory parameter (pH, pCO₂, pO₂, HCO₃) scores between experimental and control group was statistically significant. It indicates that there was significant difference between the groups. Based on posttest pH, pCO₂, pO₂, HCO₃ mean scores, experimental group showed improvement in all respiratory parameters compared with control group.

The Paired t' test value of cholinesterase scores showed statistical significance in experimental group (t=4.00). Whereas control group showed statistical non significance (t=0.71). Unpaired t' test value of posttest cholinesterase scores between experimental and control group was statistically significant (t=3.84). It indicates that there was significant difference between the groups. Based on posttest cholinesterase mean scores, experimental group (Mean=2.13) showed increased level of cholinesterase compared with control group (Mean=1.27).

This finding was supported by Murat S, (2003), concluded that organophosphorous poisoning is a serious condition that needs rapid diagnosis and treatment. Since the respiratory failure is the major reason for mortality, careful monitoring, appropriate management and nursing care, early recognition of complication may decrease the mortality rate among organophosphorous poisoning patient. There was significant inhibition of plasma cholinesterase and elevation of serum amylase at admission with return to normal values on 5th day. Plasma cholinesterase

inhibition <10% is associated with high degree of mortality. Hyperamylasemia >200 U/L has been associated with poor prognosis and proneness to respiratory failure Sharan Badiger et al, (2016).

References

- [1] Eddleston M, (2000). Patterns and problems of deliberate self-poisoning in the developing world. *Q J Med.* (93):715-31.
- [2] Eddleston M & Phillips MR, (2004). Self poisoning with pesticides. *BMJ.* (328):42-44.
- [3] Buckley NA & Karalliedde L, et al, (2004). Where is the evidence for the management of pesticide poisoning - is clinical toxicology fiddling while the developing world burns? *J Toxicol Clin Toxicol.* (42):113-6.
- [4] Darren M Roberts & Cynthia K Aaron, (2007). Managing acute Organophosphorus pesticide poisoning. *BMJ.* (334):629-34.
- [5] Kummer W & Lips KS, et al. (2008). The epithelial cholinergic system of the airways. *Histochem Cell Biol.* (130):219-234.
- [6] Takemura Y & Helms MN, et al. (2013). Cholinergic regulation of epithelial sodium channels in rat alveolar type 2 epithelial cells. *Am J Physiol Lung Cell Mol Physiol.* (304):428-437.
- [7] Akgur S & Veral A, et al.(2008). Adult respiratory distress syndrome in human organophosphate poisoning cases. *Toxicol Environ Chem* (90):493-499.
- [8] Dudekula Moulali & D.V.H.S Sarma, et al. (2014). A rare incidence of spontaneous coronary artery dissection in a case of chronic cocaine abuse - a case report. *International Journal of Recent Trends in Science and Technology Science and Technology.* 12(1): 102-104
- [9] Karalliedde L. (2002). Cholinesterase estimations revisited: the clinical relevance. *Eur J Anaesthesiol.* 19(5):313-6
- [10] Amala Rajan & Ilavarasi Jesudoss, (2016). Organophosphate poisoning: A case report, overview of management and nursing interventions. *CMI.* 14(1) :40-45.
- [11] Getnet Mequanint Adinew & Assefa Belay Asrie, (2017). Pattern of acute organophosphorus poisoning at University of Gondar Teaching Hospital, Northwest Ethiopia. *BMC Research Notes.* (10):149
- [12] Subhash Chandra Joshi & Chandra Prakash, (2013). *J Indian Acad Forensic Med.* Vol. 35, (4): 346-48.
- [13] Murat S. Intensive Care management of organophosphorus insecticide poisoning, *Indian Journal of Critical Care;* June 6, 2003; Pp. 259-260.
- [14] Sharan Badiger et al, (2016). Study of serum amylase & serum cholinesterase in organophosphorus compound poisoning. *JKIMSU.* 5(2): 49-55.